



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME

New names and the final members of new combinations are in **heavier type**

- Acacia*, 475, 476, 480
Acaciaphyllites, 420
Acarospora sp., 293
Acer californicum, 336; *rubrum*, 225, 227-229, 231
Achillea lanulosa, 237, 239
Aconitum columbianum, 237; *Lycotium*, 313
Acrostichum citrifolium, 173
Actinella, 16; *lanata*, 239
Adiantum pedatum, 54
Adopogon, 12, 16
Adoxa moschatalina, 237
Aecidium alliicola, 375
Aeodes nitidissima, 511, 512
Agoseris, 16; *acuminata*, 17; *attenuata*, 17; *altissima*, 17; *agrestis*, 17; *arachnoidea*, 18, 21; *aurantiaca*, 18, 20, 237; *elongata*, 19; *glaucia*, 17, 18, 21; *gracilens*, 20; *gracilentia*, 19; *gracilentia* Greenei, 19; *graminifolia*, 19, 20; *grandiflora*, 20, 21; *Greeneana*, 19; *Greenei*, 18, 19; *humilis*, 19; *laciniata*, 18, 21; *Leontodon*, 18; *maculata*, 17; *major*, 19; *obtusifolia*, 20; *parviflora*, 20; *pubescens*, 17; *pumila*, 17; *roseata*, 17; *rostrata*, 19, 20; *scorzoneraefolia*, 17; *tenuifolia*, 20; *turbinata*, 20; *villosa*, 17, 18
Agrimonia Bicknellii, 450; *gryposepala*, 449; *hirsuta*, 449; *mollis* **Bicknellii**, 450
Agropyrum Scribneri, 237
Alcea americana, 94; *indica*, 94
Alectoria bicolor, 291; *divergens*, 291; *jubata implexa*, 291; *ochroleuca*, 292; *sarmentosa*, 292
Algae of Lower California, Mexico, Some marine, 489
Algites, 420; **americana**, 401
ALLARD, H. A. Some common species of *Crataegus* at Thompson's Mills, Georgia, 25
Allium, 374; *acuminatum*, 374; *brevistylum*, 373, 374; *pikeanum*, 237, 241; *reticulatum*, 373; *validum*, 374
Alnus rhombifolia, 335, 336, 341, 342
Alsine baicalensis, 239; *crassifolia*, 239, 240
Alsinopsis obtusiloba, 237; *propinqua*, 237
Alsophila capensis, 546; *decussata*, 548, 550
Alternaria, 135, 137
Althaea, 94
Amarella plebeja, 238, 241
Amelanchier alnifolia, 335, 337; *canadensis*, 453; **nantucketense**, 453-455; *oblongifolia*, 453, 455; *oblongifolia micro-petala*, 454; *rotundifolia*, 455; *sanguinea*, 455; *spicata*, 454, 455
Amnicola Yatesiana, 331
Amphicosmia multiflora, 547, 548
Amygdalaceae, 337
Amygdalus Persica, 460
Ananthacorus, 158, 159, 171-173, 190; *angustifolius*, 172, 176, 181, 188
Anatheca dichotoma, 502, 503, 513; *furcata*, 503
Anchistea virginica, 55
ANDREWS, F. M. Conjugation of two different species of *Spirogyra*, 299
Andromeda, 420, 421, 423; *grandifolia*, 420; *Novae-caesareae*, 420; *Parlatorii*, 420
Andropogon virginicus, 225, 231
Androsace carinata, 237; *subumbellata*, 237
Androvettia, 404
Anetium, 153, 156-159, 171, 173; *citrifolium*, 157, 173, 188
Angelica Grayi, 237
Anodonta cygnea impura, 331
Antennaria, 382; *aprica*, 237; *media*, 237, 241; *microphylla*, 237, 239; *nardina*, 237; *rosea*, 237, 239
Anthoceros levis, 221
Anticlea elegans, 239
Antrophyopsis, 175; *immersum*, 176
Antrophyum, 158, 159, 164-183, 187-190; *anetioides*, 169; *Boryanum*, 172; *discoideum*, 169; *Dussianum*, 169; *ensiforme*, 169; *Jenmani*, 169; *latifolium*, 156; *lineatum*, 171; *minimum*, 164; *nanum*, 175, 179, 188; *plantagineum*, 172, 174, 189; *reticulatum*, 169, 172, 176, 177, 181, 182, 190; *semicostatum*, 174, 175, 188; *stenophyllum*, 189; *Werckleanum*, 164; *Williamsi*, 190
Aphanolejeunea, 251, 254, 255, 271-273, *gemmae* of, 279-286; **crenata**, 276, 277, 286; **exigua**, 273-277, *gemmae* of, 281-285; **microscopica**, 273, 275, *gemmae* of, 281-284; **sicaefolia**, 277-279, 286
Aquilegia coerulea, 237; *saximontana*, 237
Arabis Drummondii, 237, 239
Aralia, 415; *anisiloba*, 417; *groen-*

- landica, 408; spinosa, 516; washingtoniana, 409; Wellingtoniana, 415
Araliopsis, 413; **breviloba**, 417; **cretacea**, 413, 417; **cretacea dentata**, 416; **cretacea salisburyae**, 416
Araucaria, 405, 420, 421; *bladenensis*, 405, 420; *Jeffreyi*, 420
Araucarites ovatus, 405
Arbutus, Dioecism in the trailing, with notes on the morphology of the seed, 531
Arbutus Menziesii, 335, 339, 342
Archilejeunea Sellowiana, 219; *viridis-sima*, 219
Arctostaphylos, 341; *Manzanita*, 335, 339, 340
Arenaria Fendleri, 237; *sajanensis*, 237
Argentina, 351; *Anserina*, 352-354, 447; *argentea*, 352, 354; *Babcockiana*, 353-355; *Egedii*, 353-356; *litoralis*, 352-355, 447, 448; *occidentalis*, 352-354; *pacifica*, 353; *subarctica*, 352, 353, 355
Aristida stricta, 231
Aronia arbutifolia, 452; *atropurpurea*, 452; *nigra*, 452
Artemisia Pattersonii, 237; *scopulorum*, 237
Arthonia patellulata alnicola, 287
Arthoniaceae, 287
ARTHUR, J. C. New species of *Uredineae*—VIII, 369
Arundinaria macrosperma, 227; *tecta*, 225, 226
Arundo, 422; *groenlandica*, 420
Asclepiodora viridis, 232
Aspergillus niger, 135, 137
Aspidiophyllum trilobatum, 410
Asplenium cecilensis, 403, 424; *Dicksonianum*, 409; *ebenoides*, 55; *montanum*, 56; *pinnatifidum*, 77; *platyneuron*, 57; *platyneuron Hortonae*, 58; *Ruta-muraria*, 57; *Trichomanes*, 58
Assimilation of atmospheric nitrogen by fungi, Upon, 135
Aster, 437; *carolinianus*, 232
Astrodapsis Whitneyi, 340
Athyrium acrostichoides, 58; *angustifolium*, 77; *Filix-femina*, 59
Aureae, 79
Baccharis, 229; *glomeruliflora*, 372; *halimifolia*, 229; *oaxacana*, 371; *sordescens*, 371; *thesioides*, 370
Baeomyces, 287; *aeruginosus*, 288
Baptisia perfoliata, 234; *villosa*, 232
BARTLETT, H. H. A supplemental description of *Euphorbia ephedromorpha*, 343
Batodendron arboreum, 516
BENEDICT, R. C. The genera of the fern tribe *Vittarieae*: their external morphology, venation, and relationships, 153
Benzoin melissaefolium, 232
Besseyia alpina, 239
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain—VII, 399
Betulaceae, 336
Biatora cinnabarina, 288; *franciscana*, 288; *sphaeroides*, 288
BICKNELL, E. P. The ferns and flowering plants of Nantucket, 103, 447
Biflorae, 80
Bistorta bistortoides, 237; 239, *vivipara*, 237, 239
Blackberries, 103; half high, 426; high, 426
BLANCHARD, W. H. *Rubus* of eastern North America, 425
Blastenia ferruginea, 290
Botrychium dissectum, 59; *lanceolatum*, 59; *neglectum*, 59; *obliquum*, 60, 76; *simplex*, 60; *tenebrosus*, 61; *virginianum*, 61
Boykinia Jamesii, 237
Brachiolejeunea bahamensis, 205, 218; *corticalis*, 218
Brachyphyllum macrocarpum, 422
BRAINERD, E. Further notes on the stemless violets of the South, 1; The caulescent violets of the southeastern United States, 191
Brassica, 478
Brevifoliae, 81
Brunnichia cirrhosa, 233
Bryum coronatum, 34
Buellia papillata, 287; *scabrosa*, 287
Caesalpinia, 423
Calamagrostis, 319; *purpurascens*, 237
Callicostella pallida, 36
Callymenia Requienii, 511
Caltha leptosepala, 237, 239, 240
Calycites, 420; *diospyrififormis*, 418
Campanula rotundifolia, 237; *uni-flora*, 237, 239
Camptosorus rhizophyllus, 61
Candicans, 82
Canna flaccida, 232
Cannabinus, 94
Carduus scopulorum, 237, 239
Carex 422, 482; *alpina*, 237; *capillaris*, 237; *chalciolepis*, 237; *Clarkii*, 420; *ebenea*, 237; *festiva*, 237; *lupulina*, 481; *lupulina albomarginata*, 482; *melanocephala*, 237; *petasata*, 237; *rupestris*, 237; *scopulorum*, 237; *Walteriana*, 226
Carex lupulina, A new variety of, 481
Carinifex Sanctae-clarae, 331
Carnegiea, 473, 475, 476, 477; *gigantea*, 474

- Castilleja lancifolia, 237; occidentalis, 237; sulphurea, 237, 239
 Caudalejeunea Lehmanniana, 219
 Caulinites, 402; inquirendus, 401
 Caulescent violets of the southeastern United States, The, 191; synopsis of, 197
 Celastrus, 537; arctica, 407; scandens, 537
 Centroceras clavulatum, 492, 509; cryptacanthum, 509; cryptacanthum longiar-ticulatum, 509; oxyacanthum, 509
 Celastrophylloids, 421, 423; crenatum, 420; elegans, 420
 Cephalanthus occidentalis, 335, 339
 Cephalotaxospermum carolinianum, 420
 Ceramium clavulatum, 509
 Cerastium beeringianum, 237; oreophi-lum, 237
 Ceratiola ericoides, 225, 234
 Ceratolejeunea cubensis, 213, 215; **in-tegrifolia**, 213, 215, 222; patentissima, 214
 Cercocarpus, 341; betulaeifolius, 335, 337
 Cetraria arctica, 292; cucullata, 292; islandica, 292; juniperina pinastri, 292; nivalis, 292; Oakesiana, 292; Richard-sonii, 292
 Chamaecyparis thyoides, 225, 235
 Chamaenerion angustifolium, 237, 239, 241
 Chamaerhodos, 367; erecta, 367; erecta parviflora, 367; Nuttallii, 367
 Cheilanthes lanosa, 62
 Cheilolejeunea decidua, 212; phyllobola, 210, 212; versifolia, 208
 Chionophila Jamesii, 237
 Chlorophyceae, 490
 Cholisma ferruginea, 232, 516
 Chondria obtusa paniculata, 508
 Chondrophylla americana, 237, 239
 Cinnamomum, 420, 421, 423; inter-medium, 423; **Newberryi**, 420, 423
 Cissites, 415, 417; Heerii, 415; Harkeri-anus, 414, 415; obtusum, 416; salis-buriaefolius, 414-417
 Cissus laciniata, 477
 Citrophylloids, 423; aligerum, 420
 Cladium effusum, 231, 235
 Cladonia, 287; belliflora coccocephala, 288; cariosa, 289; cariosa Dawsoniana, 289; cariosa majuscula, 289; coccifera, 288; coccifera pleurota, 288; crispata, 288; crispata subcrispata, 288; digitata, 288; furcata scabriuscula, 288; pyxidata neglecta, 288; rangiferina, 288; rangi-formis, 288; sp., 292; uncinatis, 288
 Cladoniaceae, 288
 Cladophlebis, 403, 409; Browniana, 409; **socialis**, 409
 Cladophora Hutchinsiae, 491; **Mac-Dougallii**, 491, 492, 513; ovoidea, 491, 492; trichotoma, 492
 Claytonia megarrhiza, 237
 Clementsia rhodantha, 237
 Clethra alnifolia, 229
 Cliftonia monophylla, 231, 233
 Codium, 494; **decorticatum**, 494, 495; elongatum, 494, 495; galeatum, 493; Lindenbergii, 494; tomentosum, 493-495; tomentosum proliferum, 494
 Coilodesme-?, 496; californica, 496
 Collema melaenum, 290; melaenum poly-carpum, 290
 Collemaceae, 290
 Cololejeunea, 251-261, gemmae of, 279-286; Biddlecomiae, 254, 256, gemmae of, 280, 281; calcarea, 251, 252, 254, 256, gemmae of, 280, 281; ceatocarpa, 265; ciliatilobula, 254; cuneifolia, 253; diaphana, 259, gemmae of, 282, 284, 285; erigens, 263; Goebeli, 265; hetero-morpha, 257; Jooriana, 207, 270; lanciloba, 253, 268; longistylis, 282; Macounii, gemmae of, 281-283; mar-ginata, 262; microscopica, 273; minu-tissima, 258, 259, gemmae of, 282, 283, 285; **myriocarpa**, 256, 258, 259, gemmae of, 281-283, 285; ovalifolia, 281; scabri-flora, 263
 Colpomenia tuberculata, 495; sinuosa tuberculata, 495
 Colutea obovata, 407; primordialis, 407
 Comarum, 81, 356; angustifolium, 357; palustre, 356; tomentosum, 357
 Conferva trichotoma, 492
 Conioselinum scopulorum, 237, 239
 Conjugation of two different species of Spirogyra, 299
 Contributions to the history and biblio-graphy of the roselle, 91
 Contributions to the Mesozoic flora of the Atlantic coastal plain—VII, 399
 Cordaites, 406
 Cornaceae, 338
 Corneocyclas compressa, 331
 Cornus, 408; **cecilensis**, 408, 424; florida, 516; glabrata, 335, 338, 342
 Corydalis nobilis, 312; ochroleuca, 313
 Crataegus, 25-32, 437-458; Beadlei, 25, 29-32; **Bicknellii**, 243, 244, 455-459; chrysocarpa, 244, 455, 457; collina, 25, 29, 32; Crus-galli, 25, 26, 32, 456; Jonesae, 243; macrosperma, 458; prui-nosa, 455, 457, 458; rotundifolia, 457; rotundifolia Bicknellii, 244; **schizo-phylla**, 243, 455; spathulata, 25, 27, 32; tomentosa punctata, 459; uniflora, 25, 26, 28; Williamsii, 243
 Credneria rhomboidea, 413
 Crepidispermum, 16
 Crepis, 12, 15; acuminata, 14; alpicola, 13; ambigua, 15, 16; angustata, 13; barbiger, 14; biennis, 13; denticulata, 13; exilis, 14; glauca, 12, 13; glauclla,

- 12, 13; gracilis, 13, 14; intermedia, 13; occidentalis, 14; occidentalis costata, 14; occidentalis gracilis, 13; perplexans, 12, 13; petiolata, 12, 13; platyphylla, 12, 13; pumila, 14; riparia, 12, 13; riparia parva, 13; runcinata, 12; runcinata alpicola, 13; runcinata hispidulosa, 13; scopulorum, 13; scopulosa, 13; **seselifolia**, 14; tomentulosa, 12, 13
- Crossotolejeunea, 252
- Croton, 423
- Crotonophyllum, 420, 423
- Cryptomeric inheritance in *Onagra*, 461
- Cunninghamites elegans, 422
- Cyathea, 545; arborea, 550; capensis, 545; horrida, 545; multiflora, 545, 547, 548, 550
- Cyathea multiflora, type of the genus Hemitelia R. Br., On the identity of, 545
- Cyclodictyon Liebmanni, 36
- Cyclolejeunea, gemmae of, 279; accedens, 274
- Cypripedium Reginae, 232
- Dalbergia, 407; **severnensis**, 407, 424
- Dammara borealis, 400; cliffwoodensis, 400
- Dasiphora, 356, 362; fruticosa, 237, 238
- Dasystephana Parryi, 238; Romanzovii, 238
- Datura, 541; laevis, 541
- Delphinium, 313; azureum, 308; decorum scaposum, 307; nudicaule, 308; scaposum, 307-310, 312, 317
- Dennstaedtia punctilobula, 62
- Dentaria diphylla, 192
- Deschampsia alpicola, 237; caespitosa alpina, 237
- Description of two new fossil figs from Wyoming and Montana, 389
- Development of the embryo sac and endosperm in some seedless persimmons, 379
- Dewalquea, 422; Smithi, 420
- Dewberries, 426
- Dicranella Hilariana, 33; Liebmanniana, 33; mexicana, 33; Perrotettii, 33; tenuirostris, 33; trematodontifolia, 33
- Dictyota Bartayresiana, 498; Binghamiae, 499; crenulata, 499; **Vivesii**, 497-499, 512
- Diocism in the trailing arbutus, with notes on the morphology of the seed, 531
- Diospyros, 418, 421, 423; brachysepala, 418; primaeva, 417, 418, 420; rotundifolia, 420; **vera**, 418; virginiana, 418, 379; virginiana Turneri, 339
- Diplasiolejeunea, 255, 262, 267; pellucida, 262; Rudolphiana, 207; unidentata, 207
- Dodecatheon pauciflorum, 237, 239
- Dodonaea, 295
- Doryanthites**, 406; **cretacea**, 406
- Draba aurea, 238; Parryi, 238; streptocarpa, 238
- Drepanolejeunea Araucariae, 276; infundibulata, 274
- Dryas octopetala, 238
- Drymocallis, 80, 86, 362; agrimonioides, 362-367; albida, 362, 365; amplifolia, 362, 363; arguta, 362, 366, 367; arizonica, 362, 363; ashlandica, 366, 367; convallaria, 362, 366, 367; corymbosa, 362; cuneifolia, 366; fissa, 363, 366, 367; foliosa, 362, 363; glabrata, 363, 366, 367; glandulosa, 363, 364, 367; glandulosa incisa, 362; glandulosa monticola, 362; glutinosa, 362, 366, 367; gracilis, 366; Hanseni, 363, 366; incisa, 366; lactea, 366; laxiflora, 362, 364, 365; micropetala, 362, 365; oregana, 362, 363, 365; pseudorupestris, 366, 367; pumila, 362, 363; reflexa, 364-366; rhomboidea, 363, 366, 367; valida, 362, 363, 367; viscida, 365, 367; viscosa, 362, 363; Wrangelliana, 363-367
- Dryopteris Boottii, 63; Clintoniana, 77; cristata, 63, 64; cristata $\times$ intermedia, 63; cristata $\times$ marginalis, 64; Goldiana, 65, 76; intermedia, 65, 66; intermedia $\times$ marginalis, 65; marginalis, 66; noveboracensis, 66; simulata, 77; spinulosa, 66; Thelypteris, 66
- Duchesnea, 357; indica, 357
- Early spring aspects of the coastal plain vegetation of South Carolina, Georgia, and northeastern Florida, 223
- Echinocactus, 477
- EGGLESTON, W. W. New Crataegi of the northeastern manual range, 243
- Eichhornia crassipes, 382
- Elephantella groenlandica, 238, 239
- Enteromorpha compressa, 491; intestinalis, 490; Linza, 490
- Ephebe pubescens, 290
- Epidendrum conopseum, 516
- Epigaea, 531-542; dioecism in, 535; evidence of heterostyly in, 534; morphology of the seed, 538; repens, 531, 533, 537, 538, 540, 541, 543
- Epilobium anagallidifolium, 238; ovatifolium, 238; stramineum, 238, 239
- Equisetum arvense, 73; fluviatile, 73; hyemale, 73; hyemale affine, 74; sylvaticum, 74
- Erianthus sp., 225, 226
- Ericaceae, 339
- Ericoideae, 540
- Erigeron elatior, 238; formosissimus, 238, 239; leucotrichus, 241; pinnatisectus, 238; simplex, 238, 239, 241; Smithii, 238, 239; uniflorus, 241

- Eriocaulon decangulare*, 229
Eritrichium argenteum, 238
Erysimum oblancoletum, 238, 239
Euantrophyum, 175, 176
Eucalyptus, 334, 421, 423; *angusta*, 421; *Geinitzi*, 421; *Wardiana*, 421
Eulejeunea albida, 208; *flava*, 207
Euonymus europaeus, 537
Euosmolejeunea clausa, 212; *duriuscula*, 212; *opaca*, 212; *trifaria*, 213
Eupatorium rotundifolium, 229
Euphorbia ephedromorpha, 343-346
Eupolytaenium, 171
Eusargassum, 496
Euvittaria, 166
 EVANS, A. W. *Hepaticae* of Puerto Rico, 251; *The Hepaticae* of the Bahama Islands, 205
Evernia prunastri thamnoides, 292
 Evidence of recapitulation in the *Vittarieae*, 186

Fabaceae, 338
Fagaceae, 336
Faucha, 507; *laciniata*, 506; **mollis**, 507, 508, 513; *nitophylloides*, 507, 508; *repens*, 506; **Sefferi**, 506, 513
 Ferns and flowering plants of Nantucket, The, 103, 447
Festuca brachyphylla, 238; *ovina ingrata*, 238
Ficus, 389-392, 400, 420; *atavina*, 421; *Bruesi*, 391; *callicarpa*, 391; **Ceratops**, 389; *crassipes*, 421; *elastica*, 400; *Krausiana*, 421; **matawanensis**, 399, 424; *neurocarpa*, 391; *ovata*, 410; *ovatifolia*, 410; *protogaea*, 391; *punctata*, 391; *reticulata*, 400; **Russelli**, 392; *Stephensoni*, 421; *Woolsoni*, 410
 Figs from Wyoming and Montana, Description of two new fossil, 389
Filix bulbifera, 67; *fragilis*, 67, 238
Fissidens polypodioides, 33
Flabellaria, 405
 Flora of Lehigh County, Pennsylvania, 45
Fragaria, 357; *americana*, 358, 359, 361; *bracteata*, 358; *californica*, 357; *canadensis*, 358; *chiloensis*, 360, 361; *cuneifolia*, 360; *firma*, 359; *glauca*, 238-240, 359, 360; *grandiflora*, 357, 360, 361; *Grayana*, 360, 361; *illinoensis*, 360; *insularis*, 359; *mexicana*, 357; *multicipita*, 357, 358; *ovalis*, 359; *pauciflora*, 359; *platypetala*, 358, 359; *prolifera*, 359; *pumila*, 359; *Suksdorfii*, 359, 360; *Terrae-novae*, 358; *vesca*, 357-361, 447; *virginiana*, 357-361, 447; *yukonensis*, 359, 360
Frasera carolinensis, 232
Fraxinus oregona, 336
Frigidae, 79

Frullania, 205, 256; *arietina*, 220; *caroliniana*, 220; *gibbosa*, 220; *obcordata*, 220; *riojaneirensis*, 220; *squarrosa*, 220
Fucus tomentosus, 493-495
 Further notes on the stemless violets of the South, 1
Fusarium, 135, 137; *Zeae*, 135

 GAGER, C. S. *Cryptomeria* inheritance in *Onagra*, 461
Gelsemium sempervirens, 516
Gemmae of *Cololejeunea*, *Leptocolea*, and *Aphanolejeunea*, 279
 Genera of the fern tribe *Vittarieae*: their external morphology, venation, and relationships, The, 153
Gentiana cruciata, 312, 313; *Parryi*, 238; *plebeja*, 238, 239, 241; *Romanzovii*, 238
 Georgia, Some common species of *Crataegus* at Thompson's Mills, 25
Geum agrimonoides, 362; *canadense*, 449; *virginianum*, 449
Gigartina Chamissoi, 502; *Chauvinii*, 502; *papillata*, 502; *Teedii*, 502
Ginnania undulata, 502
Glandulosa, 80
Gleichenia gracilis, 402; *Saundersii*, 402, 424
Glossophora Kunthii, 499
Gonidea angulata Cooperi, 331
Gracilaria, 504, 508; *armata*, 504; *confervoides*, 504; *compressa*, 504; *Cunninghamii*, 504; *multipartita*, 504; *peruana*, 504, 505; **Vivesii**, 503-505, 513
Graciles, 83; *pinnatae*, 83, 85
Grammitis pumila, 162
Grossularia Menziesii, 335, 337
Grossulariaceae, 337
Gymnogramma, 163

Halimeda, 493; *discoidea*, 492, 493; *Tuna*, 493
Halymenia, 510; **actinophylla**, 509-513; *coccinea*, 511; *floridana*, 510, 511; *latifolia*, 510; *ulvoidea*, 510, 511
Hamamelites, 423; *cordatus*, 421
 HANNIBAL, H. A *Pliocene* flora from the Coast Ranges of California, 329
 HARPER, R. M. Early spring aspects of the coastal plain vegetation of South Carolina, Georgia, and northeastern Florida, 223; The relation of climax vegetation to islands and peninsulas, 515
 HARRIS, J. A. Teratological fruits of *Ptelea*, 385
Hecistopteris, 156, 157, 159, 162, 179, 182, 189; *pumila*, 162; *Werckleana*, 155, 164
Heckera umbellata, 382
Hedera, 423; *primordialis*, 421
Hemionitis brasiliana, 169; *cayennensis*,

- 169; lanceolata, 169; lineata, 169; reticulata, 174
Hemitelia, 545; *Hartii*, 548; multiflora, 546, 548, 550; *nigricans*, 547, 548, 550; *obscura*, 548, 550; *Parkeri*, 546
Hemitelia R. Br., On the identity of *Cyathea* multiflora, type of the genus, 545
Hepaticae of Puerto Rico, 251
Hepaticae of the Bahama Islands, The, 205
Heteranthera limosa, 382
Heterolepis, 420
Heteropleura, 16; *ambigua*, 15, 16; **Fendleri**, 15, 16
Heterothecium sanguinarium, 288
Heuchera Hallii, 238; *parvifolia*, 238
Hibiscus, 97; *digitatus*, 97; *gossypifolius*, 97; *Sabdariffa*, 91, 97
Hicoria, sp., 516
Hieracium, 14, 15; *albertianum*, 15; *albiflorum*, 15; *canadense*, 14, 15; *columbianum*, 14, 15; *cynoglossoides*, 15; *Fendleri*, 15; *griseum*, 15; *macranthum*, 15; *nigrocollinum*, 15; *runcinatum*, 12, 13; *scabriusculum*, 15; *Scouleri*, 15; *umbellatum*, 14, 15
Holomitrium crispulum, 33
Hordeum jubatum, 377, 378
Howe, M. A. Phycological studies—V. Some marine algae of Lower California, Mexico, 489
Howe, R. H. Jr. List of lichens collected in the Yukon region by Mr. R. S. Williams, 287
Hyophila Oerstediana, 34
Hypocoum Matthioli, 94
Hypopterygium Tamarisci, 36
Hyptis stellulata, 369
Ilex, 407; *glabra*, 226, 231; *myrtifolia*, 226, 231–233; *opaca*, 516; **severnensis**, 407, 424
Illicium, 420, 423
Indica elegans, 94
Induced and occasional parasitism, 473
Iris hexagona, 232
Isoetes, 154; *canadensis*, 76; *Engelmanni*, 76
Isopappus divaricatus, 225
Ivesia, 80

Jatropha stimulosus, 378
Juglans, 422; *arctica*, 420
Juncoides parviflorum, 238; *spicatum*, 238
Juncus castaneus, 238; *Drummondii*, 238; *Roemerianus*, 229; *triglumis*, 238
Jungermannia cuneata, 264; *flava*, 207; *marginata*, 262, 266; *microscopica*, 273
Jungermanniaceae, 206
Juniperus, 235; *barbadensis*, 229, 235; *virginiana*, 231, 235
Jussiaea grandiflora, 229

KINGSLEY, M. A. On the anomalous splitting of the rhizome and root of *Delphinium scaposum*, 307
KNOWLTON, F. H. Description of two new fossil figs from Wyoming and Montana, 389
Krameria, 479
Krigia, 12, 16; *virginica*, 12
Kuhnistera pinnata, 225, 234

Lactuca, 21, 22; *augustana*, 22, 23; *integri-
folia*, 23; **polyphylla**, 23; *saligna*, 22; *Scariola*, 22; *Scariola integrata*, 22; *virosa*, 22, 23
LAMSON-SCRIBNER, F. Notes on certain species of *Muhlenbergia*, 319
Laurencia glandulifera, 508; *paniculata*, 508; *patentiramea*, 508; *pinnatifida*, 508
Laurophyllum, 421, 423; *elegans*, 420; *nervillosum*, 420
Laurus, 421, 423; *atanensis*, 420; *canariensis*, 339; *plutonia*, 420
Lecanora calcarea, 290; *castanea*, 290; *epibrya*, 290; *lentigera*, 289; *pallida*, 289; *rugosa*, 290; *Schleicheri*, 290; *subfusca argentata*, 289; *subfusca campestris*, 289; *thamnoplaca*, 289; *ventosa*, 290
Lecanoraceae, 289
Lecidaceae, 287
Lecidea alpestris, 293; *arctica*, 293; *contigua hydrophila*, 288; *platycarpa*, 288; sp., 293
Leguminosites, 420–423; *robinifolia*, 420
Lehigh County, Pennsylvania, Flora of, 45
Lejeunea, 251–256, 261; *americana*, 207; *calcarea*, 252, 253; *cardiocalpa*, 267, 268; *cavifolia*, 252; *ceatocalpa*, 265; *ciliatilobula*, 254; *cuneata*, 264, 265; *cuneifolia*, 280; *epiphyta*, 257, 258; *erigens*, 262, 264; *flava*, 207, 208; *glaucescens*, 208; *Goebelii*, 265; *heteromorpha*, 257, 258; *heterophylla*, 273, 275; *hyalino-marginata*, 266; *Jooriana*, 270; *laevigata*, 253, 254; *liliputiana*, 279; *Macounii*, 281; *marginata*, 262, 266, 267; *microscopica*, 252, 253, 273; *micrandroecia*, 262; *Moorei*, 207; *myriocalpa*, 256–258; *obtusangula*, 215; *obliqua*, 254, 265; *ovalifolia*, *gemmae* of, 281–284; *pseudostipulata*, 256; *scabriflora*, 262, 264, 265; *sicaefolia*, 273, 277, 278; *Sintenisii*, 273, 275; *Wrightii*, 257
Lepidopilum brevipes, 36; *polytrichoides*, 36; *semilaeve*, 36
Leptasea *austromontana*, 238; *chrys-antha*, 238; *flagellaris*, 238
Leptocolea, 251–255, 260–262, 267, *gemmae* of, 279–286; *cardiocalpa*, 260, 268–271, 286; **ceatocalpa**, 265; *gemmae* of, 282–284; *cuneifolia*, 253, 280–284,

- gemmae of, 281, 282, 284; **Goebelii**, 265, 280, gemmae of, 282-285; **Jooriana**, 270, 271, gemmae of, 282, 283, 285; *laevigata*, 253; **lanciloba**, 253, 268, gemmae of, 282, 283; **longistylis**, 282, 283; *marginata*, 262, 266, 268; **micrandroecia**, 252, 262; **ovalifolia**, 281-284; **planifolia**, 265-268, 270, 286; **scabri-flora**, 262, 265, gemmae of, 282-286
- Leptogium myochroum saturninum*, 290
- Leptolejeunea elliptica*, 218
- Leucobryum giganteum*, 33
- Leucodoniopsis plicata*, 34
- Leucolejeunea xanthocarpa*, 220; *unciloba*, 219
- Leucophyllae*, 82, 84
- Leucothoe Catesbaei*, 322
- Lewisia pygmaea*, 238, 239, 241
- Lichens collected in the Yukon region by Mr. R. S. Williams, List of, 287
- Lindigia aciculata*, 35
- Liquidambar Styraciflua*, 225, 227, 516
- Liriodendron Tulipifera*, 225
- List of lichens collected in the Yukon region by Mr. R. S. Williams, 287
- List of plants collected above timber line on Pikes Peak, with altitudinal extensions and notes, 237
- Lloydia serotina*, 238
- Lopholejeunea Sagraeana*, 219
- Lorinseria areolata*, 77
- Loxogramma*, 183
- Lubie Endigi*, 95
- Lupinus diffusus*, 225, 234
- Luzula parviflora*, 238; *spicata*, 238
- Lychnis montana*, 238
- Lycopodium*, 420, 421; *annotinum*, 78; *clavatum*, 74; *complanatum*, 74, 78; *complanatum flabelliforme*, 74; *inundatum*, 75; *lucidulum*, 75; *obscurum*, 75; *palmatum*, 68; *tristachyum*, 78
- Lygodium palmatum*, 68
- MACDOUGAL, D. T. Induced and occasional parasitism, 473
- Machaeranthera aspera*, 238, 239
- MACKENSEN, B. Two new species of *Opuntia*, 141
- Macromitrium apiculatum*, 34; **flavopilosum**, 34; *mucronifolium*, 34; *ulophyllum*, 34
- Macrorhynchus glaucus laciniatus*, 18, 21; *purpureus*, 18, 19
- Magnolia*, 421, 423; *californica*, 339; *Capellinii*, 406, 420; *Fraseri*, 232; *glaucula*, 225-229, 231; *grandiflora*, 226-231, 234, 516; *Newberryi*, 420; *obtusata*, 420; *tenuifolia*, 420
- Malaceae, 337
- Malapoenna geniculata*, 232
- Malus Malus*, 452
- Malva*, 94
- Marchantia chenopoda*, 206
- Marchantiaceae, 206
- Marshallia trinervia*, 232
- Mastigolejeunea auriculata*, 219
- Matteuccia Struthiopteris*, 68
- Mayepea macrocarpa*, 145
- MAXON, W. R. On the identity of *Cyathea multiflora*, type of the genus *Hemitelia* R. Br., 545
- Melia Azederach*, 225
- Mertensia alpina*, 238, 241; *ciliata*, 238, 241; *picta*, 241
- Mesadenia sulcata*, 232
- Mesophytic climax, 517
- Meteoriopsis patula*, 35; *recurvifolia*, 35
- Metzgeriopsis*, 251, 252; *pusilla*, 251
- Micranthes rhomboidea*, 238, 240
- Microlejeunea bullata*, 208; *laetevirens*, 208; *lucens*, 208
- Mitchella repens*, 516
- Momisia*, 420, 422
- Monogramma*, 153, 156, 158-163, 167, 182, 190; *dareicarpa*, 160-163, 178, 180, 182, 185, 186, 188; *graminea*, 160-163, 166, 178, 182, 185, 187, 189; *paradoxa*, 160-162, 164, 167, 168, 178, 186, 189; *subfalcata*, 160, 161, 178, 179, 180, 186, 189; *trichoidea*, 160-162, 178, 186, 189
- Monotropoideae, 540
- Moriconia americana*, 422
- Morus alba*, 145; *celtidifolia*, 146; **mollis**, 145
- Muhlenbergia*, Notes on certain species of, 319
- Muhlenbergia californica*, 323-325; *comata*, 319; **curtifolia**, 328; *curtifolia Griffithsii*, 328; *glomerata brevifolia*, 323; *huachucana*, 325; *Lemmoni*, 324-327; *neo-mexicana*, 320-322; *Parishii*, 323; *pauciflora*, 320-322; **polycaulis**, 327; *Pringlei*, 320; sp., 377; *sylvatica californica*, 323; *sylvatica Pringlei*, 320, 321
- Multifidae, 82, 86
- Multijugae, 80, 82, 83
- Myrica*, 421, 422; *cerifera*, 226, 234, 516; *Brittoniana*, 420; *elegans*, 420
- Myrsine*, 423; *Gaudini*, 408, 420
- Nantucket, The ferns and flowering plants of, 103, 447
- Narinam-Poulli, 96
- Neckera disticha*, 35; *undulata*, 35
- Nematostylae*, 80, 356
- Nephroma arcticum*, 291; *expallidum*, 291
- New Crataegi of the northeastern manual range, 243
- New species of Uredineae, 369
- New variety of *Carex lupulina*, A. 481
- Nitophyllum ocellatum*, 508
- Nitrogen, Upon assimilation of atmospheric, by fungi, 135

- Notes on certain species of Muhlenbergia, 319
 Notes on Rosaceae, 79, 351
 Notes on Rutaceae—V. Species characters in Ptelea and Taravalia, 295
 Nymphaea sagittifolia, 232
 Nyssa biflora, 225–231; capitata, 232; Ogeche, 228, 231, 232; sylvatica, 226; uniflora, 228, 229, 233, 234
- Octoblepharum albidum, 33
 Olsynium grandiflorum, 376
 Onagra, Cryptomeric inheritance in, 461
 Onagra, 465; biennis, 461–463, 471; rubricalyx, 468; rubrinervis, 468
 Onoclea, 402, 421; inquirenda, 401, 420, 424; sensibilis, 68
 On the anomalous splitting of the rhizome and root of Delphinium scaposum, 307
 On the identity of Cyathea multiflora, type of the genus Hemitelia R. Br., 545
 Onychiopsis, 409
 Ophioglossum vulgatum, 77
 Opuntia, 474, 477, 478, 480; Blakeana, 474, 476, 478; discata, 475; **leptocarpa**, 141; Lindheimeri, 142; macrorhiza, 142, 143; **Roseana**, 142; versicolor, 477; vulgaris, 516
 Oreobroma pygmaea, 238, 240
 Oreochrysum Parryi, 238
 Oreoxis humilis, 238, 239
 Orontium aquaticum, 234
 Orthostichopsis tetragona, 35
 Osmanthus americana, 516
 Osmunda, 402; cinnamomea, 69, 229; cinnamomea frondosa, 69; Claytoniana, 69; obergiana, 401, 402; spectabilis, 69
 Oxyria digyna, 238
- Pachistima, 420, 423
 Pachypteris, 404
 Padina Durvillaei, 497; sp., 497
 Padus demissa, 335, 337
 Palmocarpus sp., 389, 390
 Paludetrina longinqua, 331
 Panama mosses, 33
 Pannaria brunnea, 291; lepidiota coralliphora, 291; pezizoides, 291
 Pannariaceae, 291
 Parkinsonia, 476
 Parmelia conspersa, 291; stygia, 291
 Parmeliaceae, 291
 Paronychia pulvinata, 238
 Pasania densiflora, 335, 336, 342
 Pecopteris socialis, 409
 Pedicularis Parryi, 238
 Pellaea atropurpurea, 69
 Peltigeraceae, 291
 Penicillium, 135, 138; glaucum, 138
- PENNINGTON, L. H. Upon assimilation of atmospheric nitrogen by fungi, 135
 Pennsylvania, Flora of Lehigh County, 45
 Pentstemon glaucus stenosepalus, 238, 239; Hallii, 238
 Persea Borbonia, 516
 Persimmons, Development of the embryo sac and endosperm in some seedless, 379
 Pertusaria dactylina, 289; rhodocarpa, 289
 Phacelia sericea, 238
 Phaeophyceae, 495
 Phalaris, 375
 Phaseolus indicus, 95
 Phegopteris Dryopteris, 77; hexagonoptera, 70; Phegopteris, 71
 Philonotis sphaericarpa, 34; nanodendra, 34
 Phleum alpinum, 238
 Phoradendron flavescens, 225, 226–231, 516
 Phragmites, 422; Pratti, 420
 Phycological studies—V. Some marine algae of Lower California, Mexico, 489
 Phyllogonium fulgens, 35; viscosum, 35
 Physa heterostrophia, 331
 Physocolea, 252
 Physostegia virginiana, 541
 Pieris nitida, 231
 Pilophorus cereolus acicularis, 289
 Pilotrichum amazonum, 35
 Pinaceae, 335
 Pinus, 422; caribaea, 519, 520; echinata, 225–229; Elliottii, 226, 231, 233, 236, 520; glabra, 227, 232, 235; palustris, 225–231, 515, 519–524; raritanensis, 421; rigida, 524; serotina, 225, 226, 228, 231; Taeda, 225–228, 231
 Piper medium, 382
 Pistia Nordenskioldi, 405, 406
 Placodium elegans, 290; ferrugineum, 290; Jungermanniae, 290; vitellinum, 290
 Plagiochila ludoviciana, 206
 Planorbis trivolvus, 331
 Platanus, 412, 415; Heerii, 409, 411–413; recurvata, 411, 412, 415
 Pleurogramma, 156, 165
 Pliocene flora from the Coast Ranges of California, A, 329
 Pluchea bifrons, 229
 Poa arctica, 238; Grayana, 238; phoenicea, 238; rupicola, 238
 Podeum, 504, 508
 Podozamites Knowltoni, 403, 421; lanceolatus, 410; marginatus, 406, 410
 Polemonium confertum, 238; pulcherrimum, 238
 Polysiphonia californica, 508

- Polystichum acrostichoides*, 71
Polypodium marginellum, 157; *poly-*
podioides, 516; *serrulatum*, 157; *vul-*
gare, 71
Polypogon, 319
Polytaenium, 159, 169–190; **anetioides**,
169; **brasilianum**, 169; **cayennense**, 169,
174, 179, 189; **discoideum**, 169, 190;
Dussianum, 169; **ensiforme**, 169, 171;
Jenmani, 169, 171; **lanceolatum**, 169,
170, 172, 176, 177, 180, 181, 190;
lanceolatum Feei, 189; *lineatum*, 169,
170, 172; **quadriseriatum**, 170–172, 178,
179, 190
Pontederia cordata, 229, 382
Populus trichocarpa, 335, 336
Porotrichum cobanense, 35
Potamogeton, 422
Porphyra leucosticta, 499
Potentilla, 79, 351; *acuminata*, 80;
alpestris, 79; *ambigens*, 85; *Anserina*,
351, 352, 354; *Anserina Egedii*, 353;
Anserina grandis, 352, 353; *Anserina*
groenlandica, 353; *Anserina sericea*,
352; *anserinoides*, 354; *arachnoidea*,
89; *argentea*, 448; *arguta*, 362; *ari-*
zonica, 86; *argyrea*, 82; *atrovirens*, 86,
89; *biflora*, 80; *bipinnatifida*, 86, 87,
238, 239; *brevifolia*, 80, 81; *Breweri*, 83;
Bruceae, 82, 83; *canadensis*, 357, 448,
449; *cascadensis*, 85; *ciliata*, 367;
coloradensis, 83; *crinita*, 84, 85; *de-*
currens, 85; *dissecta decurrens*, 85;
diversifolia pinnatisecta, 84, 85; *diffusa*,
83; *Drummondii*, 85; *Egedii*, 353;
effusa, 82, 83; *emarginata*, 79; *filipes*,
83, 238; *fissa*, 367; *flabelliformis*, 79;
fragiformis, 79; *fragiformis parviflora*,
79; *Friesiana*, 79; *glabrella*, 86, 87, 89;
glandulosa, 364, 366, 367; *glandulosa*
fissa, 366; *glandulosa genuina*, 366;
glandulosa nevadensis, 367; *glauco-*
phylla, 238; *glutinosa*, 367; *gracilis*, 83;
Hickmani, 84; *Hippiana*, 82, 83; *Hip-*
piana pulcherrima, 83; *klamathensis*,
83, 84; *Keilhavii*, 87; *lactea*, 366, 367;
lasiodonta, 86; *Lemmoni*, 84, 85;
litoralis, 86, 88; *lupina*, 82; *lupulina*, 82;
Macounii, 81; *maculata*, 79; *millefolia*,
84; *minutifolia*, 82; *missourica*, 88, 89;
monidensis, 85; *monspeliensis*, 448;
multifida, 86, 88; *nana*, 79; *Nelsoniana*,
84, 85; *nivea*, 79; *occidentalis*, 352;
oregana, 364; *Osterhoutiana*, 83; *ovina*,
84, 85; *pacifica*, 352–354; *palustris*, 80,
81, 356; *palustris villosa*, 357; *pauci-*
juga, 86; *pectinata*, 89; *pennsylvanica*,
82, 86–89; *pennsylvanica communis*,
88; *pinnatifida*, 86; *pinnatisecta*, 84, 85;
plattensis, 84, 85, 89; *propinqua*, 83, 86;
proxima, 81; *pseudosericea*, 86, 87;
pulchella, 86, 87; *pulchella elatior*, 87,
88; *pulcherrima*, 83; *pumila*, 449; *re-*
flexa, 364, 367; *reptans*, 357; *rhomboi-*
dea, 367; *Richardii*, 85; *rosulata*, 80;
rubricaulis, 81, 82; *rubripes*, 81–83, 86;
rupestris, 366; *rupestris americana*, 366;
rupicola, 83; *Salesoviana*, 356; *saxi-*
montana, 82, 238; *saxosa*, 80, 81;
sericea glabrata, 89; *simplex*, 448;
Sommerfeltii, 87; *strigosa*, 86, 89;
subarctica, 88, 353; *trichocarpa*, 80,
356; *tridentata*, 80; *Vahlana*, 87;
versicolor, 83, 84; *virgulata*, 86; *vir-*
dior, 82, 83, 238, 239; *Wahlana*, 87;
wyomingensis, 84, 85; *Wrangelliana*,
367
Prenanthes pauciflora, 12
PRETZ, H. W. *Flora of Lehigh County*,
Pennsylvania, 45
Primula angustifolia, 238; *Parryi*, 238
Prionodon densus, 35
Prionolejeunea, 252
Protea, 422
Proteoides, 420–422; *lancifolius*, 421
Protophyllocladus, 420, 421; **lobatus**, 403;
polymorphus, 404; *subintegrifolius*, 404,
405
Protophyllum multinerve, 411; *Stern-*
bergii, 411
Prunus americana, 460; *angustifolia*, 255;
Avium, 459; *Cerasus*, 459; *maritima*,
459; *serotina*, 459
Pseudocryphaea flagellifera, 34
Pseudocymopterus multifidus, 238, 239;
purpureus, 238, 239
Pseudotsuga taxifolia, 335
Psoralea physodes, 335, 338
Ptelea, 295–297; *angustifolia*, 295;
Baldwinii, 295; *monophylla*, 234; *to-*
mentosa, 297; *trifoliata*, 295–297, 385;
trifoliata mollis, 297; *viscosa*, 295
Ptelea, *Teratological fruits of*, 385
Pteridium aquilinum, 72
Pteridophyta, 54
Pteris, 403; *Albertsii*, 403; *angustifolia*,
172; *graminea*, 159; *lineata*, 164
Pterospermites, 411
Ptilocalais macrolepis, 11; *major*, 11;
nutans, 11
Ptilophora major, 11
Ptiloria, 11; *pauciflora*, 12; *ramosa*,
11, 12
Puccinia Allii, 375; *Asparagi*, 375; *Blas-*
dalei, 374, 375; *Calochorti*, 376;
egregia, 370; **exornata**, 370; **fidelis**, 369;
granulispora, 375; *mutabilis*, 373, 375;
oaxacana, 370; **pagana**, 372, 376;
pistorica, 372; *Porri*, 375; *rubigovera*,
378; *Sisyrinchii*, 376; **sphenica**, 371;
Winteriana, 375
Puerto Rico, *Hepaticae of*, 251
Pyroloideae, 540
Pyrus communis, 452

- Quercus*, 420, 422; *agrifolia*, 335, 337; *Catesbaei*, 225, 227, 231; *Chapmani*, 516; *chrysolepis*, 335-337, 342; *falcata*, 229; *geminata*, 516; *laurifolia*, 516; *marylandica*, 225; *nigra*, 225-229, 231; *Phellos*, 229; *virginiana*, 229, 516, 519
- Radiovittaria*, 166
- Radula*, 251; *australis*, 206
- Ramalina polymorpha*, 292; *pusilla*, 292; *minuscula*, 292
- Ranunculaceae*, 311
- Ranunculoides*, 79-81
- Ranunculus inamoenus*, 238, 239; *multifidus*, 382
- Relation of climax vegetation to islands and peninsulas, The, 515
- Rectolejeunea Berteroana*, 208, 210; *Brittoniae*, 209, 211, 212, 222; *flagelliformis*, 210; *phyllobola*, 211, 212
- Relationships of the Vittarieae, 183
- Rhamnaceae*, 338
- Rhamnus californica*, 335, 338, 342; *Purshiana*, 335, 338, 342
- Rhizogonium spiniforme*, 34
- Rhodiola integrifolia*, 238, 242; *polygama*, 238, 242
- Rhodophyceae*, 499
- Rhodophyllis bifida*, 508
- Rhodymenia*, 504; *peruviana*, 505
- Rhodymenioidae*, 508
- Rhus*, 423
- Rhynchosstegium scariosum*, 36
- Ribes lentum*, 238, 241
- Ricciaceae*, 206
- Ricciella crystallina*, 206
- Rivales*, 85
- Robinia hispida*, 232
- Roripa alpina*, 238, 239
- Rosa*, 437; *carolina*, 450; *cinnamomea*, 451; *humilis*, 451; *rubiginosa*, 451; *rugosa*, 451; *sp.*, 450; *virginiana*, 450
- Rosaceae*, 103, 337, 447
- Rubiaceae*, 339
- Rubricaulis*, 81
- Rubus* of eastern North America, 425
- Rubus*, 425, 429, 432, 435-439; *abbrevians*, 119; *allegheniensis*, 106, 426-431, 435, 438; *allegheniensis* × *argutus*, 111; *allegheniensis* × *frondosus*, 112, 114; *allegheniensis* × *procumbens*, 113; *amabilis*, 436; *amicalis*, 436; *Andrewsianus*, 106, 426, 429, 430, 434, 438; *angulatus*, 439; *arenicolus*, 123; *argutus*, 106, 116, 118, 429; *argutus* × *Baileyanus*, 116; *argutus* × *flagellaris*, 117, 119; *argutus* × *frondosus*, 114; *argutus* × *nigricans*, 115; *argutus* × *procumbens*, 116, 121; *ascendens*, 115; *Baileyanus*, 109, 116, 119, 122, 128, 130, 131, 438; *Baileyanus* × *Ensenlii*, 130; *Baileyanus* × *flagellaris*, 132; *Baileyanus* × *frondosus*, 118, 119, 121; *Baileyanus* × *procumbens*, 127; *biformispinus*, 120; *Boyntoni*, 438; *caesius*, 432; *canadensis*, 125, 426, 430, 432, 435; *canadensis frondosus*, 435; *canadensis Millspaughii*, 427; *cuneifolius*, 426, 434; *Ensenlii*, 109, 123, 131, 132, 438; *Ensenlii* × *flagellaris*, 131; *Ensenlii* × *frondosus*, 123; *Ensenlii* × *hispidus*, 125; *Ensenlii* × *procumbens*, 128; *flagellaris*, 108-110, 118, 119, 127, 132, 430, 438; *flagellaris* × *frondosus*, 118, 119; *flagellaris* × *hispidus*, 126; *flagellaris* × *procumbens*, 129; *floricomus*, 106; *floridus*, 438; *frondosus*, 106, 107, 119, 122, 123, 427-429, 433-435; *frondosus* × *hispidus*, 120; *frondosus montanus*, 428; *frondosus* × *nigricans*, 119; *frondosus* × *procumbens*, 121; *fruticosus*, 427, 428, 430; *geophilus*, 128, 130; *grandicaulis*, 437; *hispidus*, 108, 124, 125, 127, 426-428, 430, 432-438; *hispidus major*, 436; *hispidus* × *nigricans*, 124; *hispidus* × *procumbens*, 111, 124; *hispidus suberectus*, 436; *Millspaughii*, 426-429; *multiformis*, 437; *multispinus*, 121; *nigricans*, 107, 108, 124, 126, 436; *nigrobaccus*, 106, 427, 429; *nitidus*, 439; *obovalis*, 430; *occidentalis*, 105; *parvifolius*, 434; *philadelphicus*, 435; *procumbens*, 109, 111, 116, 121, 122, 128, 131, 426, 431, 433, 438; *procumbens roribaccus*, 127; *pubescens*, 432; *Randii*, 437; *recurvans*, 107, 426, 433; *recurvicaulis*, 437; *roribaccus*, 128, 131; *semisetosus*, 436; *sempervirens*, 430; *setosus*, 431, 435, 436; *strigosus*, 105, 238, 242; *suberectus*, 427, 429; *triflorus*, 105, 432; *trivialis*, 426, 428, 431-433, 439; *vermontanus*, 125, 436; *villosus*, 427-432, 435; *villosus Randii*, 433
- RUSBY, H. H. Two new species of edible fruits from Mexico, 145
- RYDBERG, P. A. Notes on Rosaceae, 79, 351; Studies on the Rocky Mountain flora, 11
- Rydbergia grandiflora*, 238
- Rynchospora dodecandra*, 516
- Sabal glabra*, 227, 229, 231, 235; *Palmetto*, 232, 235
- Sabalites*, 405; *magothiensis*, 405
- Sabbatia decandra*, 232
- Sabdariffa*, 92-94; *Alia*, 92; *Lobellij*, 92
- Salicaceae*, 336
- Salix*, 334, 420-422; *brachycarpa*, 238, 240, 242; *flexuosa*, 421; *fluvialis*, 335, 336; *laevigata*, 335, 336; *Lesquerexii*,

- 409, 421; nigra, 227; pseudo-Hayei, 421; saximontana, 238
- Santa Clara formation, 329
- Sapindus, 423; Morrisoni, 421
- Sargassum-?, 496; Agardhianum, 496; Filipendula, 496; Liebmanni, 496
- Sassafras, 413-415; acutilobum, 409, 414, 415; cretaceum, 409, 412-415, 417; cretaceum dentatum, 414, 416; cretaceum obtusum, 414-416; cretaceum recurvatum, 412; integrifolium, 414; mirabile, 412, 413, 415; Mudgei, 414; obtusum, 414; recurvatum, 411, 412; subintegrifolium, 414
- Saxifraga austromontana, 238; cernua, 238; chrysantha, 238; debilis, 238; flagellaris, 238; rhomboidea, 238
- Saxosae, 80
- Scinaia, 500; furcellata, 502; furcellata complanata, 500, 502; furcellata undulata, 500, 502; **latifrons**, 500, 502, 512
- Scirpus Eriophorum, 229
- Scoliosorus, 171
- Scorzonera, 16
- SCRIBNER, F. LAMSON-. Notes on certain species of Muhlenbergia, 319
- Sedum, 313; stenopetalum, 238
- Selaginella apus, 76; densa, 238, 240; rupestris, 78
- Sematophyllum pungens, 36
- Senecio atratus, 238; carthamoides, 238; chloranthus, 238, 240; crocatus, 238; eremophilus, 238, 240; Fendleri, 238; pudicus, 238; taraxicoides, 238; werneriaefolius, 238, 240
- Sequoia, 341; Reichenbachii, 421, 422; sempervirens, 335, 336, 342
- Serenoa serrulata, 231, 235, 516
- Seseli, 14
- SHERFF, E. E. A new variety of Carex lupulina, 481
- Sibbaldia, 357; procumbens, 239
- Sibbaldiopsis, 80, 362
- Sieversia turbinata, 239
- Significance of Monogramma, 185
- Silene acaulis, 239
- Sinapis, 478
- Sisyrinchium grandiflorum, 376
- Smilax auriculata, 232; lanceolata, 229; laurifolia, 225, 231; pumila, 516
- Solidago decumbens, 239
- Solorina crocea, 291; saccata spongiosa, 291
- Some common species of Crataegus at Thompson's Mills, Georgia, 25
- Sophia incisa, 239, 240
- SOTH, B. List of plants collected above timber line on Pikes Peak, with altitudinal extensions and notes, 237
- Sphaerium sp., 331
- Sphaerophoraceae, 287
- Sphaerophorus coralloides, 287; fragilis, 287
- Spiraea latifolia, 447; tomentosa, 447
- Spirogyra communis, 299; crassa, 299
- Spirogyra, Conjugation of two different species of, 299
- Stellaria crassifolia, 239, 240; umbellata, 239
- Stephanomeria, 11; tenuifolia, 12; pauciflora, 12
- Stereocaulon, 287; paschale, 289; tomentosum, 289; tomentosum simplex, 289
- Stereophyllum **Howei**, 35
- Sterigmatocystis niger, 137
- STEVENS, N. E. Dioecism in the trailing arbutus, with notes on the morphology of the seed, 531
- Sticta pulmonaria linita, 291
- Stictolejeunea, 270
- Stramonium, 94
- Strawberry, pine, 361
- Strobilites, 420
- Studies on the Rocky Mountain flora—XXV, 11
- Stylopappus grandiflorus, 18; laciniatus, 18, 20, 21; laciniatus longifolius, 21
- Subjugae, 83
- Suffruticulosae, 356
- Supplemental description of Euphorbia ephedromorpha, A, 343
- Swertia palustris, 239
- Sympetalae, 540
- Synthyris alpina, 239
- Syrrhopodon Bernoullii, 33; brachysteloides, 33
- Taravalia, 295-297; aptera, 296
- Taraxacum, 21; angustifolium, 21; dumeatorium, 21, 239, 240; leiospermum, 21
- Taxilejeunea **obtusangula**, 215, 217, 218, 222
- Taxithelium planum, 36
- Taxodium, 235; distichum, 227-231, 235; imbricarium, 225, 226, 228, 231, 235
- Telesonix Jamesii, 237
- Teratological fruits of Ptelea, 385
- Tetraneuris, 16; lanata, 239
- Thalictrum alpinum, 239, 240; purpurascens, 382
- Thelotrema lepadinum, 290
- Thinnfeldia, 403, 404; rhomboidalis, 404
- Thlaspi coloradense, 239
- Thyrsopteris, 409
- Tillandsia tenuifolia, 516; usneoides, 225-227, 229, 231, 234, 516, 518
- Tonestus pygmaeus, 239
- Tragopogon Dandelion, 16; lanatum, 16; virginicum, 16
- Trautvetteria carolinensis, 232
- Trichostelium fluviale, 36
- Trifolium dasyphyllum, 239; nanum, 239

- Trilisa odoratissima*, 226
Trionum, 95
Trisetum subspicatum, 239
Troximon, 16, 18; *arachnoideum*, 18, 21; *arizonicum*, 19; *aurantiacum*, 19, 237; *elatum*, 19; *glaucum*, 17; *glaucum dasycephalum*, 17; *glaucum laciniatum*, 18, 21; *gracilens*, 18, 19; *gracilens Greenei*, 19; *grandiflorum*, 18, 20, 21; *grandiflorum laciniatum*, 21; *grandiflorum obtusifolium*, 20, 21; *grandiflorum tenuifolium*, 20, 21; *laciniatum*, 17, 18, 20; *montanum*, 20; *pumilum*, 17; *purpureum*, 18, 19; *roseum*, 17; *villosum*, 17
Trypethelium cruentum, 516
 Two new species of edible fruits from Mexico, 145
 Two new species of *Opuntia*, 141
Ulmus sp., 227
Ulva decorticata, 494; *fasciata*, 490; *Lactuca*, 490; *Lactuca rigida*, 490; *Linza*, 490; *rigida*, 490
Umbilicaria hyperborea, 291; *Muhlenbergii alpina*, 291; *pustulata*, 291
Umbilicariaceae, 291
 Upon assimilation of atmospheric nitrogen by fungi, 135
Urceolaria scruposa, 290
Uredineae, New species of, 369
Uromyces aemulus, 373, 375; *agnatus*, 378; *ambiguus*, 374, 376; *aterrimus*, 374; *aureus*, 374, 375; *bicolor*, 374, 376; *Hordei*, 377; *Houstoniatus*, 376; *Jatrophae*, 378; *major*, 377; *minus*, 377; *mysticus*, 377; *oaxacanus*, 378; *primaverilis*, 376; *probus*, 376; *reticulatus*, 376; *Sisyrrinchii*, 376
Usnea plicata, 292
Usneaceae, 291
Vaccinium vacillans, 433
Vaccinioideae, 540
Valvata virens, 331
Vaseya comata, 319
Veronica alpina, 239, 240
Viburnum, 339
Viola affinis, 1, 2; *amorphophylla*, 3, 4; *arvensis*, 197, 198; *asarifolia*, 195; *blanda*, 7-9; *Brainerdii*, 8; *Brittoniana*, 5, 6, 9; *canadensis*, 1, 191, 194, 197; *chalcosperma*, 2; *clandestina*, 195; *conspersa*, 195-197; *cucullata*, 2, 3, 7; *cucullata* × *fimbriatula*, 3, 4; *debilis*, 196; *dentata*, 4, 5; *denticulosa*, 7; *Egglestonii*, 9; *emarginata*, 4, 5, 7; *eriocarpa*, 194, 197; *esculenta*, 2, 9; *fimbriatula*, 2-5, 7, 194; *fimbriatula* × *palmata*, 3; *hastata*, 191-193, 197; *hastata glaberrima*, 192; *hirsutula*, 192; *incognita*, 7-9; *incognita Forbesii*, 8; *insignis*, 6; *labradorica*, 195; *lanceolata*, 7; *Langloisii*, 1; *Langloisii pedatiloba*, 2; *lobata*, 192; *multicaulis*, 196; *odorata*, 196; *pallens*, 7; *palmata*, 3, 7; *pedata*, 6, 9; *pedatifida*, 6, 7, 9; *Porteriana*, 4, 5; *primulifolia*, 5, 7; *primulifolia australis*, 7; *primulifolia villosa*, 7; *pubescens*, 1, 194, 197; *punctata*, 195, 196; *Rafinesquii*, 197; *renifolia*, 7, 8; *repens*, 195; *rostrata*, 196, 197; *rotundifolia*, 195; *sagittata*, 2, 4, 5, 7, 194; *sagittata emarginata*, 5; *scabriuscula*, 1, 194; *septemloba*, 5-7, 9; *sororia*, 195; *striata*, 192-194, 196, 197; *tenuipes*, 193; *tripartita*, 191, 192, 197; *tripartita glaberrima*, 192-194, 197; *uliginosa*, 195; *viarum*, 2, 7; *villosa*, 2, 7, 192; *vittata*, 7; *Walteri*, 192, 196, 197
Vitis rotundifolia, 516
Vittaria, 154-190; *elongata*, 189; *Gardneriana*, 166; *intramarginalis*, 165, 170, 176, 178, 180, 187, 189, 190; *lineata*, 165, 166, 177, 178; *minima*, 164-166, 178, 188; *minor*, 165; *Orbignyana*, 166; *pusilla*, 165; *remota*, 163, 166, 176-178, 180, 183, 187-190, 275; *scolopendrina*, 189; *scabrida*, 188; *sikkimensis*, 164, 178-180, 189; *stipitata*, 166, 188
Vittarieae, 153, 158; genera, 158; general considerations, 182; ontogenetic stages of, 176; pluriseriate genera of, 167; tribal characters, 155
Vittarieae: their external morphology, venation, and relationships, The genera of the fern tribe, 153
 WESTER, P. J. Contributions to the history and bibliography of the roselle, 91
Widdringtonites subtilis, 421, 422
 WILLIAMS, R. S. Panama mosses, 33
 WILSON, P. Notes on Rutaceae—V. Species characters in *Ptelea* and *Taravalia*, 295
 WOODBURN, W. L. Development of the embryo sac and endosperm in some seedless persimmons, 379
Woodsia ilvensis, 72; *obtusa*, 73
 Yukon region, List of lichens collected in the, 287
Zizania aquatica, 227, 229
Zizia cordata, 232
Zygadenus elegans, 239